



ARTICLE

WILEY

Constructions of infinite graphs with Ramsey property

Péter Komjáth

Institute of Mathematics, Eötvös
University, Budapest, Hungary

Correspondence

Péter Komjáth, Institute of Mathematics,
Eötvös University, Pázmány P. s. 1/C,
1117, Budapest, Hungary.
Email: kope@cs.elte.hu

Funding information

Thematic Excellence Programme,
Industry and Digitalization
Subprogramme, NRDI office, 2019

Abstract

For every infinite cardinal λ and $2 \leq n < \omega$ there is a directed graph D of size λ such that D does not contain directed circuits of length $\leq n$ and if its vertices are colored with $< \lambda$ colors, then there is a monochromatic directed circuit of length $n + 1$. For every infinite cardinal λ and finite graph X there is a λ -sized graph Y such that if the vertices of Y are colored with $< \lambda$ colors, then there is a monochromatic induced copy of X . Further, Y does not contain larger cliques or shorter odd circuits than X . The constructions are using variants of Specker-type graphs.

KEYWORDS

directed graphs, infinite graphs, Ramsey property, Specker type graphs, undirected graphs

1 | INTRODUCTION

A. Joó proved that if $n \geq 2$, κ is an infinite cardinal, then there is a directed graph (V, D) with no directed circuit of length at most n , such that if V is colored with κ colors, then there is a monochromatic directed circuit of length $n + 1$ [3]. The cardinality of V is 2^κ and Joó asked if there is a similar example with $|V| = \kappa^+$. In this note we give such an example using a variant of Specker-type graphs.

With a slight modification of the method we prove another result. If X is a finite graph and λ is an infinite cardinal, then there is a graph Y such that $|Y| = \lambda$ and $Y \Rightarrow (X)_{<\lambda}^1$. That is, if $\gamma < \lambda$ and the vertices of Y are colored with γ colors, then there is a monocolored, induced copy of X . Further, Y does not contain larger cliques or shorter odd circuits than X . If X does not contain odd circuits, that is, if it is bipartite, then for every $1 \leq s < \omega$ we can construct a Y with

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Journal of Graph Theory* published by Wiley Periodicals, Inc.

$Y \Rightarrow (X)_{<\lambda}^1$ and $\text{oddgirth}(Y) = 2s + 1$ by applying the same result to the vertex disjoint union of X and a circuit of length $2s + 1$.

The same method gives the existence of a hypergraph $\mathcal{H} \subseteq [\lambda]^n$ such that the chromatic number of \mathcal{H} is λ and if $H \neq H' \in \mathcal{H}$, $|H \cap H'| \geq 2$, then the elements of $H \cap H'$ are in the same positions in H and H' , that is, if $y \in H \cap H'$, then $y = x_i^H = x_i^{H'}$ for some $i < n$, where $H = \{x_j^H : j < n\}_{<}$, $H' = \{x_j^{H'} : j < n\}_{<}$ are the monotonic enumerations of H, H' .

The fact that if X is a finite graph and κ is an infinite cardinal, then there exists a graph Y with $Y \Rightarrow (X)_\kappa^1$ is well known and easy to prove. One possibility is the following. If the vertex set of X is $V = \{v_0, \dots, v_k\}$, $3 \leq k < \omega$, $\lambda = \exp_{k-1}(\kappa)^+$ we define Y as follows. The vertex set is $[\lambda]^k$, and if $A = \{a_0, \dots, a_k\}_{<} \in [\lambda]^{k+1}$, then join $A - \{a_i\}$ and $A - \{a_j\}$ in Y iff $\{v_i, v_j\} \in X$. If $F : [\lambda]^k \rightarrow \kappa$ is a vertex coloring of Y , then by the Erdős-Rado theorem there is a set $S \in [\lambda]^{k+1}$ such that all k -element subsets of S get the same color. If now $S = \{s_0, \dots, s_k\}$, $B_i = S - \{s_i\}$, then by construction $\{B_i, B_j\} \in Y$ iff $\{v_i, v_j\} \in X$, that is, $[S]^k$ gives an induced monochromatic copy of X (see [2] and [4]).

Another construction gives that if μ, κ are cardinals, $\mu \leq \kappa$, κ infinite, then for every graph X on μ , there is a graph Y with $|Y| = 2^\kappa$, with $Y \Rightarrow (X)_\kappa^1$. The vertex set of Y is ${}^\kappa\mu$, if $f, g \in {}^\kappa\mu$, then $\{f, g\} \in Y$ iff for $\delta = \min \{\alpha < \kappa : f(\alpha) \neq g(\alpha)\}$ one has $\{f(\delta), g(\delta)\} \in X$. A Baire category-type argument shows that $Y \rightarrow (X)_\kappa^1$. (A similar idea is used in Joó's above mentioned construction).

The following generalization is presented in [5]. There, if n, t are finite, κ infinite, X is an n -chromatic graph, $|X| \leq \kappa$, a graph Y is given with $Y \rightarrow (X)_\kappa^1$ such that $|Y| = \kappa^+$, and each subgraph of Y of size $< t$ is n -colorable. Notice that here only a not necessarily induced monochromatic copy of X is guaranteed.

The advantage of our construction over these examples is that it gives an induced copy and it has optimal size: κ^+ as opposed to $\exp_{k-1}(\kappa)^+$ or 2^κ .

In the definition of the directed and undirected graphs we utilize Specker graphs, that is, when the underlying set is $[\lambda]^t$ for some finite t , with $\bar{x}, \bar{y} \in [\lambda]^t$ joined if $\bar{x} \cap \bar{y} = \emptyset$ and they interlace in a predetermined way.

Specker-type graphs were introduced by Specker in [7] originally to prove the negative ordinary partition relation $\omega^3 \not\rightarrow (\omega^3, 3)^2$. Then Erdős and Hajnal used them in [1] to give examples of large chromatic graphs with arbitrarily large (finite) odd girth. Our method differs from those in [1] and [5] in that not one but finitely many interlacing possibilities are allowed.

1.1 | Notation

Definition We use the notions and notation of axiomatic set theory. In particular each cardinal is the least ordinal with that cardinality. If κ is a cardinal, $\exp(\kappa) = \exp_1(\kappa) = 2^\kappa$, and recursively $\exp_{r+1}(\kappa) = \exp(\exp_r(\kappa))$.

If f is a function, A a set, then $f[A] = \{f(x) : x \in A\}$. If S is a set, κ a cardinal, then $[S]^\kappa = \{\bar{x} \subseteq S : |\bar{x}| = \kappa\}$. Finite subsets of some set are typically denoted by \bar{x}, \bar{y}, \dots . The notation $\{x_0, x_1, \dots, x_{n-1}\}_{<}$ means that $x_0 < x_1 < \dots < x_{n-1}$.

A partially ordered set $(P, <)$ is a pair where $<$ is a binary relation on P which is irreflexive, transitive but not necessarily trichotomic. If $x \in P$, then $x \downarrow = \{y \in P : y < x\}$.

A graph is a pair (V, X) where $X \subseteq [V]^2$. A directed graph is a graph in which each edge has a direction, that is, $X \subseteq V \times V$, with no loops, that is, edges of the type $\langle v, v \rangle$. If X is a graph, then $\langle v_0, v_1, \dots, v_n \rangle$ is a cycle, if $\{v_i, v_{i+1}\} \in X$ for $i < n$, where formally $v_n = v_0$. Here, n is the length of the

cycle. A *circuit* is a cycle with distinct vertices. $\text{oddgirth}(X)$ is the length of the shortest circuit of odd length. Similarly, in a directed graph a *directed cycle* is $\langle v_0, \dots, v_n = v_0 \rangle$ where $\langle v_i, v_{i+1} \rangle \in D$ ($i < n$). A *directed circuit* is a directed cycle with distinct vertices.

A *clique* is a complete subgraph in some graph. The *clique number* is the size of the largest clique (if finite).

If X, Y are graphs, γ a cardinal, then $Y \Rightarrow (X)_\gamma^1$ denotes that whenever the vertex set of Y is colored with γ colors, then there is a monocolored induced copy of X . $Y \Rightarrow (X)_{<\kappa}^1$ denotes that $Y \Rightarrow (X)_\gamma^1$ holds for every $\gamma < \kappa$. If a not necessarily induced copy is guaranteed, then we write $Y \rightarrow (X)_\gamma^1$.

A *hypergraph* \mathcal{H} on V is any set of subsets of V , where $1 < |H| < \omega$ holds for $H \in \mathcal{H}$. Its *chromatic number*, $\text{Chr}(\mathcal{H})$ is the least cardinal κ such that there is a *good coloring* $f: V \rightarrow \kappa$, that is, such that $|f[H]| \geq 2$ ($H \in \mathcal{H}$).

Definition If $F: [\kappa^+]^t \rightarrow \kappa$, $\gamma < \kappa$, we let $\phi_t^\gamma(x_0, \dots, x_{t-1})$ be the following formula: $x_0 < \dots < x_{t-1} < \kappa^+$ and $F(x_0, \dots, x_{t-1}) = \gamma$.

By reverse recursion on $i < t$ we define $\phi_i^\gamma(x_0, \dots, x_{i-1})$ as

$$(\exists^* x_i) \phi_{i+1}^\gamma(x_0, \dots, x_i),$$

where \exists^* is the quantifier “there exist unboundedly many.” This way, ϕ_0^γ is the formula

$$(\exists^* x_0)(\exists^* x_1) \dots (\exists^* x_{t-1})[(x_0 < \dots < x_{t-1}) \wedge (F(x_0, \dots, x_{t-1}) = \gamma)].$$

Theorem 1. If $F: [\kappa^+]^t \rightarrow \kappa$, then ϕ_0^γ holds for some $\gamma < \kappa$.

Proof. We show that for every $s \in [\kappa^+]^{\leq t}$ there is $\gamma(s) < \kappa$ such that $\phi_{|s|}^{\gamma(s)}(s)$ holds. This we prove by reverse induction on $|s|$.

If $|s| = t$, $\gamma(s) = F(s)$ applies.

If $|s| < t$ and the statement holds for all s' with $|s'| > |s|$, for every $\max(s) < \xi < \kappa^+$ we have a $\gamma(s \cup \{\xi\})$. There is $i < \kappa$ such that $\{\xi : \gamma(s \cup \{\xi\}) = i\}$ is cofinal in κ^+ . Set $\gamma(s) = i$.

The statement for $s = \emptyset$ gives the Theorem. \square

Theorem 2. If $M < \omega$, $\mathcal{H} \subseteq [M]^t$ is a partition of M , $F: [\kappa^+]^t \rightarrow \kappa$, then there is $\gamma < \kappa$, and an order preserving $g: M \rightarrow \kappa^+$ such that $F(g[H]) = \gamma(H \in \mathcal{H})$.

Proof. Let $\gamma < \kappa$ be such that ϕ_0^γ holds (cf. Theorem 1). We define $g(j)$ by recursion on $j < M$ such that

- (a) $g(j-1) < g(j)$ ($j < M$), and
- (b) if $H = \{x_0^H, \dots, x_{t-1}^H\}_{<}$ is the unique element of \mathcal{H} containing j , $j = x_i^H$, then $\phi_{i+1}^\gamma(g(x_0^H), \dots, g(x_i^H))$ holds.

The selection of $g(j)$ is possible, as by the inductive assumption the statement $\phi_i^\gamma(g(x_0^H), \dots, g(x_{i-1}^H))$ holds, therefore, there exists an appropriate $g(x_i^H) > g(j-1)$.

Having finished the construction, as $\phi_t^\gamma(g(x_0^H), \dots, g(x_{t-1}^H))$ holds, we have $F(g[H]) = \gamma$ ($H \in \mathcal{H}$) \square

Definition If $2 \leq t < \omega$, $0 \leq a \leq t$, $\bar{x}, \bar{y} \in [\kappa^+]^t$, then $R^a(\bar{x}, \bar{y})$, if $\bar{x} = \{x_0, \dots, x_{t-1}\}_{<}$, $\bar{y} = \{y_0, \dots, y_{t-1}\}_{<}$, and

$$x_a < y_0 < x_{a+1} < y_1 < x_{a+2} < y_2 < \dots < x_{t-1} < y_{t-a-1}.$$

Lemma 3. If $R^a(\bar{x}, \bar{y})$ and $g: \bar{x} \cup \bar{y} \rightarrow \text{ORD}$ is strictly increasing, then $R^a(g[\bar{x}], g[\bar{y}])$ holds.

Proof. As $x_i < y_j$ iff $g(x_i) < g(y_j)$, where $\bar{x} = \{x_i : i < t\}_{<}$ and $\bar{y} = \{y_j : j < t\}_{<}$. \square

2 | CONSTRUCTION OF A DIRECTED GRAPH

Definition For $2 \leq n < \omega$, $t = 2N + 1$, $0 < a < b = na < \omega$ with the values of N , a to be determined later, we define the directed graph D on $[\kappa^+]^t$ as follows. If $\bar{x}, \bar{y} \in [\kappa^+]^t$, $\min(\bar{x}) < \min(\bar{y})$, then $\bar{x} \leftarrow \bar{y}$ if $R^a(\bar{x}, \bar{y})$, $\bar{x} \rightarrow \bar{y}$ if $R^b(\bar{x}, \bar{y})$.

Lemma 4. D contains a directed circuit of length $n + 1$.

Proof. Define $\bar{x}^i = \{x_j^i : j \leq 2N\}$, where $x_j^i = (ia + j)T + i$ ($0 \leq i \leq n$, $0 \leq j \leq 2N$) for some $T > n$. Inspection shows that $R^a(\bar{x}^i, \bar{x}^{i+1})$ ($0 \leq i < n$) and $R^b(\bar{x}^0, \bar{x}^n)$. \square

Lemma 5. Let $\bar{x} = \{x_0, \dots, x_{2N}\}_{<}$, $\bar{y} = \{y_0, \dots, y_{2N}\}_{<} \in [\kappa^+]^{2N+1}$, $R^a(\bar{x}, \bar{y})$, $x_i < c < x_{i+k}$ with $a + 1 < i$, then $y_{i-a-1} < c < y_{i+k-a}$.

Proof. As $y_{i-a-1} < x_i < c < x_{i+k} < y_{i+k-a}$. \square

Lemma 6. Let $\bar{x} = \{x_0, \dots, x_{2N}\}_{<}$, $\bar{y} = \{y_0, \dots, y_{2N}\}_{<} \in [\kappa^+]^{2N+1}$, $R^b(\bar{y}, \bar{x})$, $x_i < c < x_{i+k}$ with $i + b + k + 1 < 2N$, then $y_{i+b} < c < y_{i+k+b+1}$.

Proof. As $y_{i+b} < x_i < c < x_{i+k} < y_{i+k+b+1}$. \square

Lemma 7. D does not contain directed circuits of length $\leq n$ (for appropriately chosen values of N and a).

Proof. Assume for a contradiction that $\bar{x}^0, \dots, \bar{x}^m = \bar{x}^0$ form a directed circuit of length $m \leq n$. Let r, s be such that we have r times $R^a(\bar{x}^i, \bar{x}^{i+1})$ and s times $R^b(\bar{x}^{i+1}, \bar{x}^i)$. Clearly $r + s = m$. Notice that $r \geq 1$ and $s \geq 1$ as otherwise we have an increasing or decreasing directed path.

Pick $c = x_N^0$, the middle element of \bar{x}^0 . Then obviously $x_{N-1}^0 < c < x_{N+1}^0$. Going around the circuit, applications of Lemmas 5 and 6 give

$$x_{N-1-r(a+1)+sb}^0 < c < x_{N+1-ra+s(b+1)}^0,$$

assuming the condition for the Lemmas hold, for which $b + 1 < N - (b + 1)(n - 1)$, that is, $N > (b + 1)n$, is more than enough.

Comparing the above two inequalities, we must have

$$N - 1 - r(a + 1) + sb \leq N - 1$$

and

$$N + 1 - ra + s(b + 1) \geq N + 1$$

that is,

$$-r(a + 1) + sb \leq 0$$

and

$$-ra + s(b + 1) \geq 0.$$

As $b = na$, we obtain $-r(a + 1) + sna \leq 0 \leq -ra + s(na + 1)$. This can be written as $-s \leq -ra + sna \leq r$, that is, $|(sn - r)a| \leq \max(r, s) < n$. If now $a = n$, then this can be rewritten as $|sn - r| < 1$, that is, $r = sn \geq n$, contradiction. \square

Theorem 8. *If λ is an infinite cardinal, $2 \leq n < \omega$, then there is a directed graph D such that D does not contain directed circuits of length $\leq n$ and if X is colored with $< \lambda$ colors, then there is, in some color class, a directed circuit of length $n + 1$.*

Proof. The case $\lambda = \omega$ follows from the existence of finite examples (see, eg, [1]). Assume therefore, that $\lambda > \omega$. It suffices to show the result for $\lambda = \kappa^+$, as otherwise, that is, if λ is a limit cardinal, we can take the vertex disjoint union of smaller examples.

For $\lambda = \kappa^+$, we let D be the directed graph investigated above. Notice the conditions on the parameters we earlier established: $a = n$, $b = na = n^2$, $N \geq (b + 1)n + 1 = (n^2 + 1)n + 1$, $t = 2N + 1$, and the directed graph is on the set $[\kappa^+]^t$.

Lemma 7 gives that there are no directed circuits of length $\leq n$. If $F: [\kappa^+]^t \rightarrow \kappa$, then, by Theorem 2, there is a copy of the directed circuit of Lemma 4 in $F^{-1}(\gamma)$ for some $\gamma < \kappa$. \square

3 | CONSTRUCTION OF AN UNDIRECTED GRAPH

In the following part of the paper we prove a similar partition result for undirected graphs.

Lemma 9. *If $R^a(\bar{x}, \bar{y})$, $R^b(\bar{y}, \bar{z})$, and $R^c(\bar{x}, \bar{z})$ then $c = a + b$ or $a + b + 1$.*

Proof. As $x_{a+b} < y_b < z_0 < y_{b+1} < x_{a+b+2}$. \square

Definition Let X be an undirected graph on $n = \{0, 1, \dots, n-1\}$, κ an infinite cardinal. Assume that $a_0 < a_1 < \dots < a_{n-1}$, $a_i = a^{i+1}$, are natural numbers with a to be determined later, $t \geq 2a_{n-1}$. We define the graph Y on $[\kappa^+]^t$ as follows. If $\bar{x}, \bar{y} \in [\kappa^+]^t$, then $\{\bar{x}, \bar{y}\} \in Y$ iff there is an edge $\{i, j\} \in X$ such that $R^{|a_i - a_j|}(\bar{x}, \bar{y})$ holds. Naturally, we need that the following holds: if $a_{i_0} - a_{j_0} = a_{i_1} - a_{j_1}$, then $i_0 = i_1, j_0 = j_1$. This is implied by the next Lemma.

Lemma 10. Assume that $a_i = a^{i+1} (i = 0, 1, \dots, n-1)$, k_i is integer, $|k_i| < a (i < n)$, $k_0 a_0 + k_1 a_1 + \dots + k_{n-1} a_{n-1} = 0$, then $k_0 = k_1 = \dots = k_{n-1} = 0$.

Proof. Assume for a contradiction that $k_0 a_0 + \dots + k_{n-1} a_{n-1} = 0$ yet not all of k_0, \dots, k_{n-1} are zero. Let r be least that $k_r \neq 0$. Then $k_r a_r + k_{r+1} a_{r+1} + \dots + k_{n-1} a_{n-1} = 0$. Dividing by a^{r+1} and reordering we obtain

$$k_r = -(k_{r+1} a + \dots + k_{n-1} a^{n-r-1}).$$

Now the RHS is divisible by a , therefore, as $-a < k_r < a$, one must have $k_r = 0$, contradicting the assumptions. \square

Lemma 11. Assume that $a_i = a^{i+1} (i = 0, \dots, n-1)$, k_i is an integer, and $|k_0 a_0 + \dots + k_{n-1} a_{n-1}| < a$, then $k_0 a_0 + k_1 a_1 + \dots + k_{n-1} a_{n-1} = 0$.

Proof. As $k_0 a_0 + \dots + k_{n-1} a_{n-1}$ is divisible by a , and has absolute value $< a$, we have $k_0 a_0 + \dots + k_{n-1} a_{n-1} = 0$. \square

Lemma 12. Y contains an induced subgraph isomorphic to X .

Proof. Set $\bar{x}^i = \{x(i, j) : j < t\} (i < n)$, where $x(i, j) = (a_i + j)n + i (i < n, j < t)$. Then $x(i_0, j_0) \neq x(i_1, j_1)$ if $\langle i_0, j_0 \rangle \neq \langle i_1, j_1 \rangle$. Further, $x(i_0, j_0) < x(i_1, j_1)$ iff either $i_0 = i_1$ and $j_0 < j_1$, or $i_0 < i_1$ and $a_{i_0} + j_0 \leq a_{i_1} + j_1$, that is, $(a_{i_0} - a_{i_1}) + j_0 \leq j_1$ or $i_1 < i_0$ and $a_{i_0} - a_{i_1} < j_1 - j_0$.

This implies that $R^b(\bar{x}^{i_0}, \bar{x}^{i_1})$ holds ($i_0 < i_1$) iff $b = a_{i_1} - a_{i_0}$. The definition of Y gives that $\{\bar{x}^{i_0}, \bar{x}^{i_1}\} \in Y$ iff $\{i_0, i_1\} \in X$. \square

Lemma 13. If $\{\bar{x}, \bar{y}, \bar{z}\}$ is a triangle in Y , $\min(\bar{x}) < \min(\bar{y}) < \min(\bar{z})$, $a \geq 4$, then there are i, j, k , such that $R^{a_j - a_i}(\bar{x}, \bar{y})$, $R^{a_k - a_i}(\bar{x}, \bar{z})$, $R^{a_k - a_j}(\bar{y}, \bar{z})$, and $\{i, j, k\}$ is a triangle in X .

Proof. By the definition of Y , if $\{\bar{x}, \bar{y}, \bar{z}\}$ is a triangle, then there are $\alpha > \beta, \gamma > \delta$, and $i > j$ such that $R^{a_\alpha - a_\beta}(\bar{x}, \bar{y})$, $R^{a_\gamma - a_\delta}(\bar{y}, \bar{z})$, and $R^{a_i - a_j}(\bar{x}, \bar{z})$. By Lemma 9, this implies that either

$$a_\alpha - a_\beta + a_\gamma - a_\delta = a_i - a_j$$

or else

$$a_\alpha - a_\beta + a_\gamma - a_\delta + 1 = a_i - a_j.$$

By Lemma 11, the latter cannot hold, so we have the former.

By Lemma 10, if we reorder this to one side, all coefficients must be 0.

Case 1. $\alpha = \gamma$.

In this case, after reordering all terms to the LHS, the coefficient of a_α is ≥ 1 , a contradiction.

Case 2. $\alpha > \gamma$.

In this case, a_α is the term with the largest index on the LHS, therefore $\alpha = i$. Subtracting the common term from both sides, we obtain

$$-a_\beta + a_\gamma - a_\delta = -a_j.$$

By the property in Lemma 9, we have the following possibilities.

Case 2.1. $\beta = j$.

Then $\gamma = \delta$, impossible.

Case 2.2. $j = \delta$.

Then $a_\beta = a_\gamma$, that is, $\beta = \gamma$.

Case 3. $\alpha < \gamma$.

We proceed as in Case 2. □

Lemma 14. *If $\bar{x}^0, \bar{x}^1, \dots, \bar{x}^{k-1}$ form a clique in Y with $\min(\bar{x}^0) < \min(\bar{x}^1) < \dots < \min(\bar{x}^{k-1})$ ($k \geq 3$), then there are distinct $i(0), \dots, i(k-1)$, such that $R^{a_{i(j_1)} - a_{i(j_0)}}(\bar{x}^{j_0}, \bar{x}^{j_1})(j_0 < j_1)$.*

Here, and later, if $a < 0$, then $R^a(\bar{x}, \bar{y})$ means $R^{-a}(\bar{y}, \bar{x})$.

Proof. By induction on k . The case $k = 3$ is covered in Lemma 11

Assume we have the statement for $\bar{x}^0, \dots, \bar{x}^{k-1}$ and $R^{a_{j_1} - a_{j_0}}(\bar{x}^{j_0}, \bar{x}^{j_1})$ for $j_0 < j_1 < k$. By Lemma 13 for any $j < j' < k$ there is $i(j, j')$ such that $R^{a_{i(j)} - a_{i(j')}}(\bar{x}^j, \bar{x}^{j'})$ and $R^{a_{i(j')} - a_{i(j)}}(\bar{x}^{j'}, \bar{x}^j)$. For any other j'' with $j < j'' < k$ we have $a_{i(j)} - a_{i(j, j')} = a_{i(j)} - a_{i(j, j'')}$ and so $i(j, j') = i(j, j'')$ ($j'' \neq j, j'$) and similarly $i(j, j') = i(j^*, j')$ ($j^* \neq j, j'$). This can only be if $i(j, j')$ is the same for any $j < j' < k$ and this will be $i(k)$. □

Lemma 15. $\text{oddgirth}(Y) = \text{oddgirth}(X)$.

Proof. Assume that $\langle \bar{x}^0, \bar{x}^1, \dots, \bar{x}^{2v+1} = \bar{x}^0 \rangle$ is a circuit in Y of length $2v+1 < \text{oddgirth}(X)$. There are $i(k) \neq j(k) < n$ such that $R^{a_{i(k)} - a_{j(k)}}(\bar{x}^k, \bar{x}^{k+1})$.

Let $t = 2N+1$ and let c be the middle element of \bar{x}^0 , that is, $c = x_N^0$. Then $x_{N-1}^0 < c < x_{N+1}^0$ and repeated applications of Lemmas 5 and 6 give

$$x_{N+T-(2v+2)}^0 < c < x_{N+T+(2v+2)}^0,$$

where

$$T = (a_{i(0)} - a_{j(0)}) + (a_{i(1)} - a_{j(1)}) + \dots + (a_{i(2v)} - a_{j(2v)}).$$

These give $|T| \leq 2v+2$. If $2v+2 < \text{oddgirth}(X) \leq a$, then $T = 0$ by Lemma 11.

Lemma 10 gives

$$\sum_{i(u)=\ell} 1 - \sum_{j(u)=\ell} 1 = 0$$

for any $\ell < n$. For each $u \leq 2v$ take a directed edge $e(u)$ from $i(u)$ to $j(u)$. This way, we obtain a directed multigraph Z whose edges, without the direction, are edges of X . Further, $d_{\text{in}}(u) = d_{\text{out}}(u)$ holds for every $u < n$ in Z . Then Z splits into the edge-union of directed circuits in X , one of them, say C , must be of odd length. But then, the undirected version of C is an odd circuit in X , a contradiction, as $2v+2 < \text{oddgirth}(X)$. \square

Theorem 16. *If X is a finite graph, $\lambda \geq \omega$ is a cardinal, then there is a graph Y of cardinality λ with the same clique number and the same odd girth as X , and $Y \Rightarrow (X)_{\gamma}^1 (\gamma < \lambda)$. This holds for edge-colored graphs as well: if $g: X \rightarrow r$ is a coloring of the edges of X , then there is an edge-coloring g' of Y such that if the vertices of Y are colored with $< \lambda$ colors then there is a monochromatic induced colored copy of X .*

Proof. Again, as in Theorem 8 it suffices to show the case $\lambda = \kappa^+$, $\gamma = \kappa$. (For $\lambda = \omega$, examples are given in [1]).

We have to select the parameters satisfying the following inequalities and equalities: $a \geq \max(4, \text{oddgirth}(X))$, $a_i = a^{i+1} (i < n)$, $N \geq a_n$, $t = 2N + 1$.

We define Y as in the Definition following Lemma 9. $Y \Rightarrow (X)_{\kappa}^1$ can be proved exactly as in the proof of Theorem 8.

The clique number and oddgirth of Y are the same as those of X by Lemmas 14 and 15.

The edge-colored version of the Theorem follows immediately from the fact that for every edge of $e \in Y$ there is a unique edge $f(e) \in X$ such that if in our arguments $g: n \rightarrow [\kappa^+]^t$ is an embedding into an induced copy of X , then $f(\{g(i), g(j)\}) = \{i, j\}$. \square

Before turning to Theorem 19, we prove two Lemmas on well founded partially ordered sets.

Lemma 17. *If $(P, <)$ is a well founded, partially ordered set, then there is a well order $<_w$ of P which extends $<$, that is, if $x < y$, then $x <_w y$.*

Proof. By recursion define the elements $p(\alpha)$ as follows. If $\{p(\beta) : \beta < \alpha\} \neq P$, we let $p(\alpha)$ be a minimal element of $P - \{p(\beta) : \beta < \alpha\}$ (exists, as $(P, <)$ is well founded). If $\{p(\beta) : \beta < \alpha\} = P$ we stop the construction. Clearly, $P = \{p(\alpha) : \alpha < \phi\}$ for an appropriate ordinal ϕ . If $x = p(\beta)$, $y = p(\alpha)$, $\beta < \alpha$, set $x <_w y$. Clearly, $<_w$ is a well ordering.

To check the other property, assume that $x = p(\beta)$, $y = p(\alpha)$, $x < y$, but $y <_w x$, that is, $\alpha < \beta$. This means, that when choosing $y = p(\alpha)$, x was as yet unselected, and so, as $x < y$, y was not minimal, a contradiction. \square

Lemma 18. *Assume that $(P, <)$ is a well founded partially ordered set, $|P| = \lambda$, λ is regular, and $x \downarrow$ has cardinality $< \lambda$ for $x \in P$. Then $<$ can be extended to a well order of type λ .*

Proof. Enumerate P as $\{p_{\alpha} : \alpha < \lambda\}$. Define

$$P_{\alpha} = \{x : \exists \beta < \alpha [x \leq p_{\beta}]\}$$

for $\alpha < \lambda$. Then $\langle P_{\alpha} : \alpha < \lambda \rangle$ is an increasing, continuous sequence of subsets with $\bigcup \{P_{\alpha} : \alpha < \lambda\} = P$, $P_0 = \emptyset$, $|P_{\alpha}| < \lambda (\alpha < \lambda)$. Let $<_{\alpha}$ be a well ordering of $P_{\alpha+1} - P_{\alpha}$ extending $<$ (Lemma 17).

Define $<_w$ on P as follows. If $x \neq y \in P$, $x \in P_{\alpha+1} - P_\alpha$, $y \in P_{\beta+1} - P_\beta$, then set $x <_w y$ if either $\alpha < \beta$ or else $\alpha = \beta$ and $x <_\alpha y$.

We show that $<_w$ is as required. It is clearly a well ordering. Assume that $x < y$. Let α be the ordinal such that $y \in P_{\alpha+1} - P_\alpha$. By construction, $x \in P_{\alpha+1}$. If $x \in P_\alpha$, then $x <_w y$. If $x \in P_{\alpha+1} - P_\alpha$, then $x <_\alpha y$ and so again $x <_w y$.

Finally, if $x \in P_\alpha$, then the initial segment of $<_w$ determined by x is a subset of P_α and $|P_\alpha| < \lambda$, so each proper segment is of size $< \lambda$, consequently the type of $<_w$ is at most λ . As $|P| = \lambda$, it is equal to λ . \square

Theorem 19. *If $\lambda > \omega$ is a cardinal, $3 \leq n < \omega$, then there is a family $\mathcal{H} \subseteq [\lambda]^n$ such that*

- (a) *if $H = \{y_0^H, \dots, y_{n-1}^H\} <_\lambda$ for $H \in \mathcal{H}$, $H, H' \in \mathcal{H}$ are such that $|H \cap H'| \geq 2$, $z \in H \cap H'$, then $z = y_i^H = y_i^{H'}$ for some $i < n$, and*
- (b) *if $F: \lambda \rightarrow \gamma$ for some $\gamma < \lambda$, then there is an $H \in \mathcal{H}$ monocolored by F .*

Proof. Again, we can assume that $\lambda = \kappa^+$.

Let t, a_0, \dots, a_{n-1} be as in the above definitions. We set $H \in \mathcal{H}$ if $H = \{\bar{x}^0, \dots, \bar{x}^{n-1}\}$ for some t -tuples such that $R^{a_j - a_i}(\bar{x}^i, \bar{x}^j)$ holds for $i < j < n$.

We have the properties required, the only problem is that the underlying set is $[\kappa^+]^t$. Define a relation $<$ on $[\kappa^+]^t$ by $\bar{x} < \bar{y}$ if $R^{a_j - a_i}(\bar{x}, \bar{y})$ holds for some $i < j < n$. The transitive closure of it is a partial ordering, again denoted by $<$. It is well founded, as $\bar{x} < \bar{y}$ implies $\min(\bar{x}) < \min(\bar{y})$. Further, if $\bar{x} < \bar{y}$, then $\bar{x} \in [\max(\bar{y})]^t$, whose cardinality is at most κ .

Using Lemma 18, we can find a well order $<_w$ of $[\kappa^+]^t$ whose order type is κ^+ and under which each $H = \{\bar{x}^0, \dots, \bar{x}^{n-1}\}$ has $\bar{x}^0 <_w \dots <_w \bar{x}^{n-1}$, and so the result is proved. \square

ACKNOWLEDGMENTS

This study was supported by Thematic Excellence Programme, Industry, and Digitalization Subprogramme, NRDI office, 2019. The author is thankful to the referees for their very careful job.

ORCID

Péter Komjáth  <http://orcid.org/0000-0002-2806-6103>

REFERENCES

1. P. Erdős and A. Hajnal, *On chromatic number of graphs and set systems*, Acta Math. Acad. Sci. Hung. **17** (1966), 61–99.
2. P. Erdős, A. Hajnal, and B. Rothchild, *On chromatic number of graphs and set-systems Cambridge Summer School in Mathematical Logic (Cambridge, 1971)*, Lecture Notes Math., **337**, Springer, Berlin, 1973, pp. 531–538.
3. A. Joó, *Uncountable dichromatic number without short directed cycles*, J. Graph Theory. (forthcoming). <https://doi.org/10.1002/jgt.22509>
4. P. Komjáth, *Ramsey-type results for metric spaces*, J. Comb. Theory A **45** (1987), 323–328.

5. P. Komjáth, *A Ramsey-style extension of a theorem of Erdős and Hajnal*, Fund. Math. **170** (2001), 119–122.
6. D. Soukup, *Orientations of graphs with uncountable chromatic number*, J. Graph Theory **88** (2018), 606–630.
7. E. Specker, *Teilmengen von mengen mit relationen*, Comment Math. Helv. **31** (1957), 302–314.

How to cite this article: Komjáth P. Constructions of infinite graphs with Ramsey property. *J Graph Theory*. 2020;1–10. <https://doi.org/10.1002/jgt.22539>